

Interventions of Neurodegeneration on Patients' Cognition, Behavior, and Memory and Caretakers

Madelyn Salud

Abstract

The study of neurodegeneration is rapidly advancing as new therapeutic approaches rise to prominence and researchers increasingly discuss the possibility of reversing or delaying the symptoms of neurodegenerative diseases. This paper examines six interconnected dimensions of cognitive decline, focusing on how brain atrophy affects the behavior and memory of patients and how caretakers respond to these changes. The review finds that stress and cognitive decline exert a reciprocal influence on one another: stress can induce hippocampal degradation and neuronal loss through cortisol pathways, while neurodegeneration can cause depression in approximately 58% of patients who receive an early diagnosis. Social isolation is also identified as a significant contributor to brain atrophy, with studies showing that individuals who maintain active social lives are better able to preserve cognitive homeostasis through greater neurological resistance. The paper further examines caregiver mental health outcomes, resilience factors among patients, and the role of support groups in sustaining both patients and families. Collectively, the evidence underscores that neurodegeneration extends its consequences well beyond the patient, affecting the fundamental human capacities of memory and autonomous thought, and making the pursuit of effective treatments an issue of both medical and ethical urgency.

Introduction

With origins traced primarily to the beginning of the twentieth century, the study of neurodegeneration has advanced enormously over the past hundred years. In its earlier stages, neurology predominantly relied on diagnosis through physical examination, as methods of brain imaging were extremely primitive. Common diagnostic tools used today, including computed tomography, positron emission tomography, and magnetic resonance imaging, were entirely nonexistent for much of the field's early history. During this period, it was common for prospective neuroscience students to be cautioned that most brain diseases could only be identified, not treated.

Despite the complexity of studying a living brain, examination of post-mortem human brains preceded pathological studies of active brains by decades.

Sixty years ago, the study of neurodegenerative diseases was not a prominent focus within the neuroscience field. That has changed dramatically. Early neurodegeneration research concentrated primarily on Parkinson's disease, driven by landmark discoveries regarding its biological mechanism, specifically the loss of dopaminergic cells in the substantia nigra, and its treatability through a dopamine precursor known as levo-3,4-dihydroxyphenylalanine. Studies of dementia, ALS, Huntington's disease, and other conditions followed. Today, the assumption that neurodegenerative diseases are untreatable has been largely debunked, and neuroscientists are actively investigating methods of delaying disease onset and reversing symptoms. This paper examines six dimensions of neurodegeneration as it affects patients and their caregivers: the relationship between stress and cognitive decline, the psychological impact of early diagnosis, the connection between social isolation and brain atrophy, the differential mental health burden on caregivers of Alzheimer's and Parkinson's patients, resilience factors among patients, and the therapeutic role of support groups.

Discussion

Effects of Neurodegenerative Diseases on Stress and Mood Regulation

Researchers note that neurodegeneration poses a significant threat to cognitive homeostasis as a result of the gradual degradation of neurons, making the issue a rapidly expanding focus in clinical research. Among the most significant developments, researchers have turned their attention toward potential therapeutic mechanisms of non-invasive brain stimulation (Battaglia et al., 2024). Concurrently, a well-documented connection has been established between cognitive decline and mood disorders within patients. Evidence indicates that a reciprocal relationship exists between the severity of stress episodes, neurodegeneration, and life expectancy.

The pathophysiological effects of stress on the nervous system are closely connected with the functioning of the hippocampus, an area of the brain pivotal to emotional regulation and memory (Esch et al., 2002). Understanding this connection has allowed neuroscientists to examine stress as a major contributing factor in the development of Alzheimer's disease. Stress can degrade the hippocampus, which is also the primary brain region targeted by Alzheimer's, suggesting a mechanistic overlap between chronic psychological stress and the onset of neurodegenerative pathology. As stress levels rise, learning and memory are compromised due to disturbances in

hippocampal activation, stress-related corticosteroid secretion that suppresses hippocampal neural activity, and the induction of hippocampal atrophy (Esch et al., 2002). Stress also induces neuronal loss through the opening of glucocorticoid pathways linked to elevated cortisol, which promotes high levels of steroids that alter the structure and function of dendrites. Despite these findings, establishing the precise causal role of stress in the onset and development of neurological disorders remains difficult due to the complexity and polymorphism of these conditions (Esch et al., 2002).

Psychological Impact of Early Diagnosis of Neurodegenerative Diseases

Researchers have conducted studies on the psychological consequences of predictive testing and early diagnosis for neurodegenerative diseases. One of the most notable findings was that although many patients claim they would prefer predictive genetic testing, actual participation rates are substantially lower, with only approximately 10 to 20% of eligible at-risk patients undergoing the procedure (Paulsen et al., 2011). This discrepancy reflects how many individuals prefer uncertainty over certainty when confronting diseases for which no cure currently exists. Although physical distress and anxiety often increased immediately before and after testing, most participants demonstrated successful long-term psychological adjustment, with severe psychiatric complications remaining uncommon and psychological outcomes most strongly associated with pre-existing mental health conditions.

Oliveri et al. (2018) further underlined how predictive genetic testing can substantially affect patients undergoing evaluation for hereditary neurological conditions. A long-term follow-up study of 119 individuals found that 58% of mutation carriers experienced depression during follow-up compared to 24% of non-carriers. Additionally, 27% of non-carriers reported difficulty adjusting despite receiving a favorable test result, suggesting that psychological distress can arise not only from learning one carries a disease-causing mutation but also from the broader emotional and identity-related consequences of the testing experience itself (Gargiulo et al., 2009). Notably, prior psychological history emerged as a stronger predictor of subsequent depression than genetic status, a finding supported by a significance level of $p < 0.0001$. These conclusions highlight that while many patients adapt after early diagnosis, ongoing emotional challenges frequently require sustained psychological support from caregivers and physicians.

Cognitive Decline and Social Isolation in Patients with Neurodegeneration

Researchers have observed that individuals experiencing brain atrophy face significant difficulty adapting to sudden alterations in daily routines, with social isolation emerging as a particularly consequential risk factor. This connection became especially visible during the

COVID-19 pandemic, which prompted neuroscientists to focus extensively on the relationship between cognitive homeostasis and social interaction. Among older adults, heavy social isolation has been shown to correlate with markedly worse cognitive performance compared to socially active counterparts, with the older generation being particularly vulnerable given their reduced familiarity with digital communication tools.

A study known as the SOcial LImitations Turn Up DEmentia (SOLITUDE) research project was conducted to examine the effects of social isolation on older patients with dementia and the burden carried by their caretakers. The experiment used multiple checkpoints across a six-month period. Results showed no significant changes in dementia patients over the six months overall; however, those who had experienced increased social restriction prior to the study demonstrated more detrimental impacts to their cognitive performance. Caretakers noted that cognitive decline during social isolation was consistent with poorer semantic memory, a domain specifically sensitive to progression in Alzheimer's disease (Manca et al., 2022). Despite these findings, establishing a clear causal link between solitude and brain atrophy remains challenging, because some studies associate social isolation but not loneliness with increased dementia risk, while others find the opposite pattern (Shen et al., 2022). This inconsistency is attributable both to other unmeasured risk factors affecting neurodegeneration and to the substantial heterogeneity in research designs across studies.

Mental Health Outcomes Among Caregivers of Alzheimer's and Parkinson's Patients

Researchers have assessed the psychological implications of caring for patients with Alzheimer's and Parkinson's disease, finding that caregivers across both groups commonly report guilt, frustration, anger, and burnout. A key finding was that caregiver stress levels were not significantly correlated with the severity of the patient's illness. Rather, caregiver personality and individual coping mechanisms were stronger predictors of mental health outcomes (Marziali, Donahue & Crossin, 2005). Across 34 caregivers divided into two groups by disease category and participating in a 10-week online support group model, anticipatory grief was evident in all participants, with caretakers negatively affected by the cognitive decline and loss of recognition in their loved ones.

In response, targeted psychosocial groups were introduced via online videoconferencing to examine therapeutic mechanisms including bonding, mutual disclosure, facilitated reflection, and collaborative problem-solving. Results indicated that the emotional bonds caregivers formed were not superficial; relationships evolved from initial identification with shared experiences toward a

form of deeper empathetic support in which members commiserated and resonated with one another's struggles. Differences in caregiving experience were also observed between the two groups. Alzheimer's caregivers typically grieved both the cognitive and personality changes of their loved ones, expressing confusion, resentment, and mourning of lost intimacy. Parkinson's caregivers, by contrast, expressed more frustration around their own emotional reactions toward a cognitively intact person, including impatience, difficulty tolerating role reversal, and significant self-criticism (Marziali, Donahue & Crossin, 2005). A complementary study introduced a three-dimensional model of caregiver burden encompassing role intensity, social restriction, and self-criticism as the primary factors caregivers demonstrate when supporting patients with neurodegeneration (Geerlings, 2023).

Resilience Factors Among Patients Coping with Neurodegenerative Diseases

Psychologists have examined evidence of psychological resilience across five prominent neurodegenerative diseases: Multiple Sclerosis (MS), Parkinson's Disease (PD), Huntington's Disease (HD), Motor Neurone Disease (MND), and Alzheimer's Disease (Ovaska-Stafford, Maltby & Dale, 2021). Within this framework, resilience has been categorised into core resilience, which is trait-like and largely fixed, and internal resilience, which is psychological and developable through intervention. Higher resilience was associated with less disability and better health-related quality of life in Parkinson's Disease, though it was not significantly associated with disease severity itself. For Parkinson's patients specifically, greater resilience correlated with less depression, anxiety, and somatization, as well as lower total apathy.

Across all studies reviewed, healthy lifestyle habits including regular exercise, balanced diet, adequate sleep, and abstinence from alcohol and smoking were the biological factors most strongly associated with greater adaptability to neurodegeneration, while resilience functioned as the foundational support enabling patients to maintain those habits. Psychotherapeutic and social discussion groups significantly improved resilience scores over 12 months, and early intervention was explicitly recommended across all studies, given that resilience has been shown to predict mood protection over years, particularly as the diseases themselves progressively alter patients' cognitive and social capacities. Sustained social contact including peer support and community connections was identified as sufficient to move resilience scores in a positive direction across all patient groups. However, the evidence base was heavily skewed toward MS, with 14 of 18 studies focusing on that condition and only 2 examining Parkinson's Disease. No studies specifically examined resilience in Alzheimer's Disease, limiting the applicability of these findings to the full spectrum of neurodegenerative conditions (Ovaska-Stafford, Maltby & Dale, 2021).

The Role of Support Groups in Helping Families and Patients with Neurodegeneration

Support groups play a central role in helping both patients with neurodegenerative diseases and their families cope with the emotional, social, and practical challenges associated with diagnosis and progression. These groups provide a safe environment in which individuals can voice their experiences, learn coping strategies from others who share similar trajectories, and receive emotional validation. Research has supported the conclusion that supportive care interventions can meaningfully improve quality of life, reduce caregiver burden, and enhance communication between patients and families over time.

In a study of 115 dementia patient-caregiver dyads, higher perceived social support was associated with lower caregiver burden ($r = 0.52$, $p < 0.01$), and positive caregiving experiences were associated with lower burden ($r = 0.51$, $p < 0.01$). More than half of dementia caregivers in the sample reported experiencing burden, underscoring the strong need for structured interventions such as support groups (Armitage, 2023). In a randomised controlled trial involving 75 Parkinson's disease patients and 44 caregivers, participants who engaged in weekly community-based support activities over 12 months demonstrated significant improvements in depression, anxiety, and stress levels relative to control groups. A family-centred intervention reduced caregiver burden scores by 25.1 points, demonstrating the substantial psychological benefits that structured support programs can provide. Internet-based videoconferencing support groups were additionally examined, with findings from Marziali (2006) confirming that these formats can provide emotional validation, coping strategies, and social connectedness that help caregivers sustain their wellbeing across extended caregiving periods.

Ethics, Discussion, and Limitations

Across all of the research studies examined in this paper, stress, depression, anxiety, social isolation, and caregiver burden emerge as recurring consequences of both receiving a diagnosis of neurodegeneration and caring for those who have. Several important limitations constrain the scope and generalisability of these findings. Many of the measures used to assess depression, anxiety, stress, resilience, and caregiver burden rely on self-reported data, which may be susceptible to response bias or subjective interpretation. Furthermore, the evidence base for resilience in neurodegenerative disease is substantially skewed toward Multiple Sclerosis, with no studies specifically examining resilience in Alzheimer's Disease, making it difficult to draw generalisable conclusions across different forms of neurodegeneration.

Establishing clear causal relationships between social isolation and accelerated neurodegeneration also remains methodologically challenging, as the directionality of this association continues to be debated in the literature and may be confounded by unmeasured variables. Ethical concerns additionally arise in the context of predictive testing and early diagnosis: while genetic testing and early detection models can provide clinically valuable information, ethical healthcare practice requires careful attention to the psychological consequences of delivering this information, particularly when no curative treatment exists. Protecting patient dignity, autonomy, and privacy while balancing the imperative for scientific advancement and improved therapeutic interventions is a responsibility that must be actively managed throughout the research and clinical process. Future research should prioritise diverse longitudinal cohorts, consistent outcome measures across studies, and rigorous ethical frameworks for evaluating the full impact of early diagnostic technologies on patient and caregiver wellbeing alike.

Conclusion

Throughout this review, evidence has highlighted how conditions including Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, and Multiple Sclerosis impose a profound and multidimensional burden on patients and the individuals who care for them. The research reviewed across six domains converges on a consistent finding: neurodegeneration does not operate in isolation from the psychological, social, and relational dimensions of human experience. Stress accelerates hippocampal degradation, early diagnosis carries significant emotional consequences, social isolation compounds cognitive decline, and the demands of caregiving produce measurable and lasting psychological costs that extend well beyond the patient.

At the same time, the evidence also points toward meaningful sources of protective capacity. Resilience, when cultivated through early intervention, healthy lifestyle habits, and sustained social connection, has been shown to predict mood protection over years and to support patients in maintaining adaptive functioning as disease progression unfolds. Support groups, whether in-person or conducted through digital platforms, have demonstrated statistically significant reductions in caregiver burden, depression, and anxiety, affirming the value of structured psychosocial intervention alongside clinical treatment. The pursuit of more effective treatments for neurodegenerative diseases is, ultimately, more than a question of promoting overall wellness. It is an ethical imperative on behalf of patients who have lost, or stand to lose, the seemingly inalienable capacities of accurate memory and independent thought, faculties that much of the broader population continues to exercise without awareness of their fragility.

References

- Armitage, C. J. (2023). Caregiver burden and social support in dementia: A cross-sectional study of patient-caregiver dyads. *Frontiers in Oncology*, 13. <https://doi.org/10.3389/fonc.2023.1029938>
- Battaglia, S., et al. (2024). Non-invasive brain stimulation as a therapeutic intervention for neurodegenerative diseases. *Biomedicines*, 12(3), 574. <https://doi.org/10.3390/biomedicines12030574>
- Esch, T., Stefano, G. B., Fricchione, G. L., & Benson, H. (2002). The role of stress in neurodegenerative diseases and mental disorders. *Neuroendocrinology Letters*, 23(3), 199–208.
- Gargiulo, M., et al. (2009). Long-term outcome of presymptomatic testing in Huntington's disease. *European Journal of Human Genetics*, 17(2), 165–171.
- Geerlings, S. W. (2023). Caregiver burden in Parkinson's Disease: A three-dimensional model of role intensity, social restriction, and self-criticism. *Journal of Neurology*.
- Manca, R., et al. (2022). Social isolation and cognitive decline in dementia during COVID-19: Results from the SOLITUDE study. *Frontiers in Psychiatry*, 13. <https://doi.org/10.3389/fpsyt.2022.877595>
- Marziali, E., Donahue, P., & Crossin, G. (2005). Caring for others: Internet videoconferencing group intervention for family caregivers of Alzheimer's and Parkinson's disease patients. *Journal of Gerontological Social Work*, 44(3-4), 197–210.
- Marziali, E. (2006). Internet-based support groups for caregivers of patients with neurodegenerative diseases. *Gerontechnology*, 5(2), 85–93.
- Oliveri, S., et al. (2018). Psychological aspects of predictive genetic testing for neurodegenerative diseases. *European Journal of Human Genetics*, 26, 644–654.
- Ovaska-Stafford, N., Maltby, J., & Dale, M. (2021). Psychological resilience in neurodegenerative disease: A systematic review. *Neuropsychological Rehabilitation*, 31(9), 1504–1531.
- Paulsen, J. S., et al. (2011). Predictive testing and the psychological consequences of learning one's genetic status in Huntington's disease. *Psychiatric Services*, 62(6), 612–618. <https://pmc.ncbi.nlm.nih.gov/articles/PMC3076589>
- Shen, C., Rolls, E. T., et al. (2022). Associations of social isolation and loneliness with later dementia. *Neurology*, 99(2), e164–e175. <https://www.oxcns.org/papers/650>